

## **1. Introduction**

Transplantation offers better survival and quality of life to those able to undergo the procedure. The widespread use of potent and specific immunosuppressive agents has significantly reduced acute cellular rejection rates but the diagnosis and treatment of Antibody Mediated Rejection (AMR) remains challenging with two different clinical phenotypes identified, clinical phenotype 1 occurs early post transplant and is associated with preformed Donor Specific Antibody's (DSA) while clinical phenotype 2 is related to denovo DSA and leads to chronic AMR(1).

More than 20,000 patients awaiting kidney transplantation in the United States are sensitized (typically owing to blood transfusion, pregnancy or previous transplants) to HLA class I and/or class II antibodies. A common approach to AMR prevention has been to avoid transplanting highly sensitized patients. However, avoiding transplant renders chronic dialysis the only option, with implications for patient survival, quality of life, as well as healthcare costs (1). Acute and chronic AMR are associated with poor outcomes after kidney transplantation. Specifically, patients with acute AMR are at greater risk for subsequent rejection, chronic AMR and graft loss. Therefore, early diagnosis and evidence based treatment is essential in order to minimize graft loss and maximize long term graft and patient outcome.

## **2. Scope**

Kidney and kidney pancreas transplant.

## **3. Recommendations, Standards and Procedural Statements**

This guideline will cover the criteria for diagnosis of clinical phenotype 1 acute AMR and its treatment. This guideline does not cover treatment of de novo Donor Specific Antibody (DSA) induced AMR occurring in the late post transplant period, de novo AMR is a different clinical phenotype and although its treatment does mirror that of acute AMR, every component of acute AMR treatment may not be used in de novo DSA induced AMR as the treatment approach is case dependant.

### **3.1 Diagnosis**

Clinically, acute rejection is defined as an acute deterioration in graft function associated with specific pathologic changes seen on transplant biopsy. Acute humoral rejection is caused by donor specific antibodies to Class I and Class II HLA antigens although other non-HLA antigens have also been recognized. The Banff criteria for classifying acute humoral rejection require the presence of (a) histological evidence of acute tissue injury e.g. acute tubular necrosis, capillaritis, tubulitis, arteritis, glomerulitis (b) presence of circulating DSA, and (c) immunologic evidence of an antibody-mediated process with positive peritubular capillary C4d staining reflecting complement activation via the classical pathway (2). We now know that the complement system can be activated by the alternative and lectin pathways and hence C4d may not always be positive in AMR (3).

Two out of three of the below criteria are required for diagnosis of AMR, this is taking in to consideration new entity of C4d –ve AMR and non-HLA induced AMR:-

- Acute microcirculatory inflammation as evidenced by [c+g+v>1] on biopsy
- DSA positive (MFI not specified, as there is recognition that circulating DSA's are likely to be lower post transplant in some cases due to absorption by the graft)
- C4d +ve by immunofluorescence or immunohistochemistry

### 3.2 Treatment

- Methylprednisolone 500mg x 3 doses, followed by 30mg prednisolone (to follow a reduction dosing regimen, reduction regimen not specified but as per treating physician/surgeon preference)
- Plasma exchange (PLEX) X7 (course can be extended if there is evidence of ongoing inflammation and AMR)
- Intravenous Immunoglobulins x 2 doses of 1g/kg of ideal body weight
- Rituximab 1g x 2 doses (given 2 weeks apart) – rituximab may not be used for all cases and decision to use will be by joint discussion between transplant surgeon and transplant nephrologist on service
- Optimisation of baseline immunosuppression

Diagrammatic representation:

Treatment	Day 1	Day 2	Day 3	Day 4	Day 5-13	Day 14-21	28 onwards
MP *	500mg IV	500mg IV	500 mg IV	Pred 30 mg od			
IVIg**	1gm/kg					1gm/kg	
Plasma exchange	-	PLEX#1	-	PLEX#2	PLEX #3-7 (on alternate days)		
Rituximab	-	-	-	-	-	1gm	1gm

\*Methylprednisolone, \*\*IVIg is to be dosed 1gm/kg of ideal body weight (max. 2gm/kg in one month), IVIg has to be approved for use by the immunoglobulin panel.

#### Special considerations:

Check HIV, hepatitis B,C and TB (quantiferon) serology prior to administration of rituximab. Consent form prior to use of rituximab.

### 3.3 DSA Monitoring:

DSA monitoring is currently done by Luminex which is a solid phase assay, Luminex was developed to be a qualitative assay and not a quantitative one and any interpretation of an antibody's Median Fluorescence Intensity should reflect this. Please check with the Head of Transplant Laboratory before requesting DSA. Antibody runs are done once per week (usually Wednesday) and a serum sample should reach the lab latest by Wednesday morning. For urgent results please contact H&I lab on ext. 4607. Below is a guide to DSA monitoring for patients that develop acute AMR

1. DSA at time of biopsy
2. After first round of PLEX (7 as per above protocol)
3. At end of AMR treatment to assess response

Subsequent DSA monitoring will be on an individual case by case basis.

### 3.4 Rationale:

#### 3.4.1 Plasmapheresis

Plasmapheresis removes circulating DSA and is used in desensitisation protocols as well as in the treatment of AMR (4-6) . Different techniques used include plasma exchange and immunoadsorption.

Whilst historic observational studies had mixed conclusions on the benefit of plasmapheresis in the treatment of AMR, one of the few RCTs in the treatment of AMR by Bohmig et al, had to be terminated due to the significant benefit seen in the interventional arm receiving immunoadsorption (4, 5, 7-10). Whilst there are recognised differences between immunoadsorption and plasma exchange, the principle of antibody removal is similar. This is shown from the published data, predominantly from desensitisation protocols on the effectiveness of rapid HLA antibody removal following plasma exchange (11).

### 3.4.2 Intravenous immunoglobulin

IVIg has numerous potential effector mechanisms which could attenuate the treatment of AMR, these include the ability to neutralise circulating DSA, inhibition of complement activation and blocking immune activation by competing for FcγRs(12) . Intravenous immunoglobulin has been shown to reduce degree of allosensitisation in highly sensitised renal patients on the transplant wait list and is frequently used in protocols for the prevention or treatment of alloimmune injury (13) . However, there is no significant evidence to show its efficacy for the indication of treating AMR when used in isolation. The only randomised control trial into the use of IVIg in allograft rejection is historic and pre-dates AMR as defined by Banff (14). In that particular study the researchers found that IVIg alone was equivalent to OKT3 in the treatment of steroid resistant rejection. Another 7 observational study published around the time, showed that the use of IVIg could improve the outcome in patients with both steroid and anti-lymphocyte antibody resistant rejection (15).

### 3.4.3 Rituximab [Truxima]

Biological agents targeting B-cells, plasma cells and their derivatives have been increasingly used as adjuvant therapy in both desensitisation and AMR treatment protocols over the past decade (4) . Rituximab and bortezomib having been the most extensively studied. Rituximab is a B-cell depleting monoclonal antibody directed against CD-20. Following its administration, rituximab depletes both immature and mature B-cells but not plasma cells. Preliminary case reports and series suggested a benefit of the addition of rituximab to the standard treatment protocols of AMR. As such, rituximab is now the commonest add on agent used in the treatment of acute AMR (4, 16). However more recently two notable studies have emerged which may cast some doubt over the benefit of rituximab for the treatment of AMR. In the RCT, Ritux ERAH, which has published its one year outcomes, analysed the effectiveness of rituximab versus placebo in 38 patients with acute AMR receiving plasmapheresis, IVIg and corticosteroids. At one year, there was no benefit of adjuvant rituximab in terms of allograft; the study was underpowered (17). The longer term outcomes have yet to be reported. This is to prescribed using the rituximab chart attached.



rituximab1g update  
jan 19.pdf

## 4. Education and Training

This guideline formalises what is now current practise in the unit and as such there is no requirement for additional training but the guideline will be distributed to all renal and transplant staff members for awareness.

## 5. Monitoring and Audit Criteria

Key Performance Indicator	Method of Assessment	Frequency	Lead
Incidence of AMR	Biopsy and information from Proton	Every 2 years	
1 year creatinine post AMR	Proton	Every 2 years	

## **6. Legal Liability Guideline Statement**

See section 6.4 of the UHL Policy for Policies for details of the Trust Legal Liability statement for Guidance documents

## **7. Supporting Documents and Key References**

### **References**

1. Djamali A, Kaufman DB, Ellis TM, Zhong W, Matas A, Samaniego M. Diagnosis and management of antibody-mediated rejection: current status and novel approaches. *Am J Transplant.* 2014;14(2):255-71.
2. Webster AC, Wu S, Tallapragada K, Park MY, Chapman JR, Carr SJ. Polyclonal and monoclonal antibodies for treating acute rejection episodes in kidney transplant recipients. *Cochrane Database Syst Rev.* 2017;7:Cd004756.
3. Loupy A, Haas M, Solez K, Racusen L, Glotz D, Seron D, et al. The Banff 2015 Kidney Meeting Report: Current Challenges in Rejection Classification and Prospects for Adopting Molecular Pathology. *Am J Transplant.* 2017;17(1):28-41.
4. Ahmed T, Senzel L. The role of therapeutic apheresis in the treatment of acute antibody-mediated kidney rejection. *J Clin Apher.* 2012;27(4):173-7.
5. Bohmig GA, Wahrmann M, Regele H, Exner M, Robl B, Derfler K, et al. Immunoabsorption in severe C4d-positive acute kidney allograft rejection: a randomized controlled trial. *Am J Transplant.* 2007;7(1):117-21.
6. Winters JL. Plasma exchange: concepts, mechanisms, and an overview of the American Society for Apheresis guidelines. *Hematology Am Soc Hematol Educ Program.* 2012;2012:7-12.
7. Brown CM, Abraham KA, O'Kelly P, Conlon PJ, Walshe JJ. Long-term experience of plasmapheresis in antibody-mediated rejection in renal transplantation. *Transplant Proc.* 2009;41(9):3690-2.
8. Lefaucheur C, Nochy D, Andrade J, Verine J, Gautreau C, Charron D, et al. Comparison of combination Plasmapheresis/IVIg/anti-CD20 versus high-dose IVIg in the treatment of antibody-mediated rejection. *Am J Transplant.* 2009;9(5):1099-107.
9. Rocha PN, Butterly DW, Greenberg A, Reddan DN, Tuttle-Newhall J, Collins BH, et al. Beneficial effect of plasmapheresis and intravenous immunoglobulin on renal allograft survival of patients with acute humoral rejection. *Transplantation.* 2003;75(9):1490-5.
10. Allen NH, Dyer P, Geoghegan T, Harris K, Lee HA, Slapak M. Plasma exchange in acute renal allograft rejection. A controlled trial. *Transplantation.* 1983;35(5):425-8.
11. Gloor JM, Winters JL, Cornell LD, Fix LA, DeGoey SR, Knauer RM, et al. Baseline donor-specific antibody levels and outcomes in positive crossmatch kidney transplantation. *Am J Transplant.* 2010;10(3):582-9.
12. Jordan SC, Toyoda M, Kahwaji J, Vo AA. Clinical aspects of intravenous immunoglobulin use in solid organ transplant recipients. *Am J Transplant.* 2011;11(2):196-202.
13. Jordan SC, Tyan D, Stablein D, McIntosh M, Rose S, Vo A, et al. Evaluation of intravenous immunoglobulin as an agent to lower allosensitization and improve transplantation in highly sensitized adult patients with end-stage renal disease: report of the NIH IG02 trial. *J Am Soc Nephrol.* 2004;15(12):3256-62.
14. Casadei DH, del CRM, Opelz G, Golberg JC, Argento JA, Greco G, et al. A randomized and prospective study comparing treatment with high-dose intravenous immunoglobulin with monoclonal antibodies for rescue of kidney grafts with steroid-resistant rejection. *Transplantation.* 2001;71(1):53-8.
15. Luke PP, Scantlebury VP, Jordan ML, Vivas CA, Hakala TR, Jain A, et al. Reversal of steroid- and anti-lymphocyte antibody-resistant rejection using intravenous immunoglobulin (IVIg) in renal transplant recipients. *Transplantation.* 2001;72(3):419-22.

16. Burton SA, Amir N, Asbury A, Lange A, Hardinger KL. Treatment of antibody-mediated rejection in renal transplant patients: a clinical practice survey. Clin Transplant. 2015;29(2):118-23.

17. Sautenet B, Blanco G, Büchler M, Morelon E, Toupance O, Barrou B, et al. One-year Results of the Effects of Rituximab on Acute Antibody-Mediated Rejection in Renal Transplantation: RITUX ERAH, a Multicenter Double-blind Randomized Placebo-controlled Trial. Transplantation. 2016;100(2):391-9.

### **8. Key Words**

Antibody mediated rejection, renal transplant, kidney pancreas transplant, rituximab, immunoglobulin, rejection.

This line signifies the end of the document

---

This table is used to track the development and approval and dissemination of the document and any changes made on revised / reviewed versions

<b>DEVELOPMENT AND APPROVAL RECORD FOR THIS DOCUMENT</b>			
<b>Author / Lead Officer:</b>	Shafi Malik		<b>Job Title:</b> Consultant Nephrologist
<b>Reviewed by:</b>	Atul Bagul, Head of Service and Consultant Transplant Surgeon		
<b>Approved by:</b>	Consultant Nephrologists, Transplant Surgeons, Renal Pharmacist and H&I Lab Director		<b>Date Approved:</b> Oct 2017
<b>REVIEW RECORD</b>			
<b>Date</b>	<b>Issue Number</b>	<b>Reviewed By</b>	<b>Description Of Changes (If Any)</b>
03/02/2021		MDT	rituximab specific drug chart add and change to Truxima
<b>DISTRIBUTION RECORD:</b>			
<b>Date</b>	<b>Name</b>	<b>Dept</b>	<b>Received</b>
31/10/2017	To all renal and transplant staff	Renal and Transplant	