



Paediatric Intensive Care Unit

Endotracheal Tube Management UHL Children’s Intensive Care Guideline

Staff relevant to:	Medical and Nursing staff caring for children in the PICU
Approval date:	September 2022
Version:	V5
Revision due:	September 2025
Written by:	Anand Patil
Trust Ref:	C116/2016

Related Guidelines and Policies:

CoMET	Procedure for Securing an Endotracheal Tube placed Orally (Comet guideline)
CoMET	Procedure for Securing an Endotracheal Tube placed Nasally (Comet guideline)
C43/2016	The Prevention of Ventilator Associated Pneumonia
C5/2014	Difficult Airway UHL Neonatal Guideline

Contents

1. Introduction	1
2. Cuffed or uncuffed ET tube	3
3. Cuff pressure monitoring	4
4. Best practices recommendations.....	4
5. Education and Training	6
6. Monitoring Compliance.....	6
7. Supporting Reference	6
8. Key Words.....	7
Appendix 1: Pre-intubation Aide Memoire & ET sizing guide	9

1. Introduction

This guideline applies to all the intubated children in the management of ETT.

Purpose:

- Decrease the incidence of unplanned extubation (UE) in PICU/CICU
- To standardize the use of cuffed endotracheal tube (ETT) in children's ICU and to facilitate the education of all staff groups
- Minimise the potential for subglottic injury secondary to inadvertent use of high cuff pressures or inappropriate sized uncuffed tubes.
- To standardize routine documentation of cuff pressures in all patients with cuffed Endotracheal tubes
- To assess the optimal state of sedation as well as readiness for extubation daily to minimize UE on the unit.

Definition:

Unplanned extubation (UE) is defined as premature removal of the ETT by the patient on mechanical ventilator (deliberate UE) or by staff during nursing and medical care (accidental extubation) ⁹. Paediatric Intensive Care Audit network group (PICANet) defines UE as a dislodgement of the ETT from the trachea, without the intention to extubate immediately and without the presence of airway competent clinical staff in the bed space, appropriately prepared for the procedure.²³

Why UE should be avoided?

UE may result in complications such as hypoxemia, and/or hypercarbia and these adverse events are associated with morbidity, prolonged mechanical ventilation, longer PICU stays and increased costs and rarely death.^{24, 25}

Repeated intubations, especially those performed emergently in a less controlled scenario increase the risk of airway injury and scarring, as well as pulmonary injury from excessive ventilation.⁷

In recent years, the rate of UE has been used to benchmark the quality of patient care in intensive care with respect to sedation, weaning, and use of restraints.¹⁰

Although the national average of UE rate is around 5 per 1000 days of invasive ventilation far from the recommended ideal rate of 1 per 100 ventilation days slowly and steadily UE rates are falling. Incidence of UE in Leicester Children's Hospital (PICU/CICU) still exceeds the recommended ideal rate with about an average of 17.6 UE per year over the last 5 years (as per morbidity data base).

The recommendations for the management of the ETT aim to decrease the UE incidence in our trust. Although cardiac arrest occurred in only 2% of events, UE in pediatric patients were associated with a 20% increased risk of life-threatening cardiovascular collapse with cardiopulmonary resuscitation required in up to 47% of patients.²⁹

Limited data is available regarding factors that contribute to UEs and subsequent reintubation of children. Despite numerous publications on UE, very few studies have assessed the preventive strategies to avoid adverse events. Most of the studies suggest significant reduction in UE rate after implementing a robust educational program that looked at identifying high risk patients, ETT care, and protocol directed sedation.¹ Methods of securing the endotracheal tube varied across studies, and the use of physical restraints yielded conflicting findings.^{2, 10}

In a multivariate analysis, risk factors for UE included

- Age < 2 years
- Male gender
- Agitation
- Inadequacy of ETT fixation
- Copious secretions as they influence stability of the ETT fixation. (Also increases risk of re-intubation in UE).
- Medical procedures / nursing cares
- Failure to assess extubation readiness
- Lack of 1:1 nursing care ²

The following factors were **NOT** found to increase the risk of UE

- Time of the day
- Intensity of work on ITU
- Lower staffing levels at night
- Nasal vs oral intubation
- Presence of restraints¹⁰

(Note: Although above factors may not cause UE independently, however they may still contribute as part of multifactorial causes of UE)

Reintubation rates:

- Reintubation rates post UE range from 14-65% ²⁸
- Factors such as prolonged periods of ventilation (>28 days), UE during care/procedure, sedation within 2 of UE, copious secretions have contributed to various extent influencing the need for reintubation.

2. Cuffed or uncuffed ET tube

Cuffed tracheal tubes are increasingly used in current practice. With an advantage of significantly reduced leakage, cuffed tubes have shown better tidal volumes during the recruitment maneuvers on both pressure and volume-based ventilation strategies which also affects the oxygenation, monitoring parameters such as airway pressures, end-tidal CO₂ and avoid risk such as aspiration.

The use of a cuffed ETT can overcome most of these issues, decreasing the need for multiple intubations, reduce costs without significant risk of adverse effects in children of all age groups ^{11, 14,}

20

Literature in support of the cuffed endotracheal tubes

Despite the longstanding use and benefits of uncuffed ETT in young children, cuffed ETTs are increasingly being used with several studies over the last 20 years demonstrating its benefits. Based on plaster molds of the pediatric airway in cadaveric studies, the funnel-shaped larynx was understood to transition towards a more cylindrical adult airway around 8 years of age. Recent CT, MRI, and video laryngoscopy studies have shown that the narrowest part of the pediatric airway is actually at the infraglottic level with an elliptical form of anteroposterior greater than transverse diameter; although, the rigid cricoid ring is still functionally the narrowest part of the larynx.³¹⁻³⁴ The newly appreciated elliptical shaped airway at the level of the cricoid cartilage is potentially vulnerable

Page 3 of 13

to damage even from a properly sealed uncuffed tube. With this, there is growing evidence that suggests uncuffed ETTs may not have better outcomes that truly offers an advantage over cuffed tubes, because the circular tube will occlude the elliptical airway only on the sides, leaving a gap in the anterior and posterior sides. The leak test done routinely to make sure that the tube is not too big, may actually show a leak, but there may be excessive pressure on the transverse walls of the trachea.^{35,36}

There is growing evidence that suggests cuffed ETT have additional clinical, environmental, and economic benefits compared to uncuffed ETT. Previously, there were concerns of tracheal mucosa damage from excessive pressures, but studies have shown they provide a reliably sealed airway at pressures ≤ 20 cm H₂O and reduce the need for tube exchanges.¹² Cuffed ETT has decreased incidences of sore throat and hoarse voice, likely from the decreased need for tube exchanges.³⁷ Studies have also shown no major difference in acute respiratory complications such as stridor after extubation between cuffed and uncuffed ETT^{12, 38} with some showing decreased postoperative complications.³⁷

In a study comparing leakage and tidal volumes in cuffed vs uncuffed ETT, cuffed ETT had significantly lower leaks and were measuring higher tidal volumes. It is assumed a majority of the leak is inspiratory due to increased intratracheal pressures. Expired tidal volume levels, which more accurately reflect true tidal volume, increased over time in the cuffed group suggesting better maintenance of lung volumes.³⁷ A retrospective study on neonates found that uncuffed ETT had a leak $>5\%$ in 75% of infants with a $>40\%$ leak occurring in 42.3% of infants around the third day of mechanical ventilation. Leaks were also associated with longer duration of mechanical ventilation.³⁹ Although cuffed ETT are more expensive than uncuffed ETT, they are environmentally and economically beneficial due to decreased tube changes and decreased medical gas consumption. In a study looking at sevoflurane consumption, they found the uncuffed group used 16.1 mL of sevoflurane compared to 6.2 mL in the cuffed group.⁴⁰

Recent Cochrane review looking at trials involving 2804 children under the age of 8 years comparing cuffed versus uncuffed ETTs, which found no difference between the groups for post-extubation stridor. However, both trials demonstrated a statistically significant lower rate of ETT exchange in the cuffed ETT group. Cochrane concluded that large RCT of high methodological quality should be conducted to help clarify the risks and benefits of cuffed ETTs for children.¹⁴ The study by Weiss et al demonstrated that the improved design of the Microcuff paediatric ETT helps to overcome important disadvantages of the use of cuffed ETTs in children.¹²

Microcuff ETT seals the trachea at cuff pressures that is generally considered safe to preserve tracheal mucosal perfusion pressure.¹³

The 'Microcuff' ETT cuff is made of ultra-thin material, which is softer and thinner. The cuff is more distally placed on the ETT shaft, which sits in trachea below the subglottis. In various studies, it has shown that the design of the Microcuff™ ETT is more age appropriate and therefore, superior to the design of most other commercially available paediatric ETTs.^{11, 20}

In studies with Microcuff ET tubes showed:

- very low rate of tube exchange rate and a very low rate of croup requiring therapy
- post-extubation stridor was similar (4%) in both cuffed and uncuffed tubes, but tracheal tube exchange rate was 2.1% in the cuffed and 30.8% in the uncuffed groups²⁰
- rate of epinephrine inhalation for postextubation croup and rate of successful extubation did not differ between cuffed and uncuffed groups. He concluded that children (<8 years) Microcuff ETT are the best cuffed ETT in critically ill children¹⁴

Shortcomings in paediatric cuffed ETT:

Many commercially available paediatric cuffed ETTs are poorly designed and that this might be one of the most important sources of morbidity associated with cuffed ETTs. ^{11, 15}

Factors that might be contributing to subglottic stenosis include ETT size, movement of the tube, duration of intubation, traumatic intubation, the presence of infection during intubation, and possibly gastroesophageal reflux.²⁰

3. Cuff pressure monitoring

When using a cuffed ETT, it is mandatory to monitor the cuff pressure. Cuff pressure is traditionally monitored every hour or at least every 12 hours. Cuff pressures should be checked after ETT position changes, change in head/neck position. Cuff pressure can be monitored using a continuous automated electronic device or intermittently by a handheld manometer. Either of the methods have been proven to be of similar efficacy.⁴¹

Animal studies have shown that airway mucosal damage can occur in as short as 15 minutes if the cuff pressure is high. Cuff pressures do change with change in head and neck position. It is therefore essential to monitor cuff pressures as part of ongoing patient safety and quality improvement initiatives ^{11, 15, 17}. Multiple studies (Khine, Deakers, Newth) using the microcuff ETT have not shown any increase in the incidence of post-extubation stridor and long-term airway sequelae. ^{18, 19} However, experienced paediatric ENT surgeons feel that significant airway injury is not always accompanied by stridor. The symptoms of airway injury may not present immediately after extubation, and symptoms might develop weeks to months after injury when silent ulcerations of the mucosa retract to cause stenosis. Only endoscopy can evidently detect all airway injuries ^{11, 16, 19, 20}. Only well planned, randomized studies using well designed, standardized tracheal tubes and performing airway endoscopy before and after extubation, may answer the question which mode of intubation might be the least traumatic in children below 8 years. For clinical practice today, airway endoscopy is warranted at least in all patients developing significant stridor after extubation or any other suspicion of airway damage. ^{11,16, 19}

4. Best practices recommendations

Structure

- Nurse-to-patient ration of 1:1
- Continuous quality-improvement program

Process

- Staff education and training
- Patient risk identification
- Standardization of procedures such as ETT fixation, stabilization, tube suctioning physiotherapy, patient hygiene, positioning, and transport
- Standardization of routes for monitoring of ETT position
- Early recognition of patients ready for extubation

- Implementation of a protocol-directed ventilation weaning plan
- Standardization of sedation practices according to sedation assessment scores
- Avoid delays in extubating in a timely fashion perhaps related to staff issues, resources and time of day
- Development of appropriate data tracking tools and data collection

Airway management:

1. Routine usage of microcuff ETT is recommended for all paediatric patients in whom high ventilator pressures are anticipated ^{3, 4}.
2. Uncuffed ETT is recommended in recent airway surgery/balloon dilatation, suspected subglottic stenosis, slide tracheoplasty patients. Consider uncuffed ETT if anticipated ventilator pressures are low, re-intubation after post-extubation stridor.
3. Consider uncuffed ETT with history of stridor in previous admission after extubation.
4. If there is no leak at a peak inspiratory pressure of 20 cm of water with cuff deflated, consider downsizing the ETT if clinical condition allows.
5. Oral versus nasal intubation
 - Oral intubation is recommended where early extubation is anticipated.
 - Nasal intubation is recommended if likely to remain ventilated for longer than 48-72hour
6. Assess daily for extubation and extubate at the earliest opportunity in safe and controlled environment.
7. ETT suctioning should be limited to the length of the ET tube in situ- Ideally ET suctioning length should be recorded on the observation chart. Oropharyngeal suction should be undertaken prior to the deflation of the cuff to minimise the risk of aspiration.
8. Aim to keep pressures less than 20cmH₂O on tracheal manometer.
 - The cuff pressure should be checked every hour unless intellicuff[®] is used and hourly monitoring is not possible, in these cases please record 12 hourly (if intellicuff[®] technology is used then record hourly).
9. Consider ENT review for recurrent stridor/upper airway obstruction post extubation
10. Continuous ETCO₂ monitoring is highly recommended
 - during endotracheal intubation and tracheostomy (all areas)
 - peri-operatively
 - for all ventilated patients in ICU
 - for all ventilated patients during inter-hospital or intra-hospital transfer

ETT length, position and fixation

1. Confirm correct placement of the ETT by chest XR(CXR)
 - CXR within last 24 hours if being transported out of the hospital⁵
2. Length of the ETT ⁶
 - Recommended length of the ETT post intubation in anaesthetic department is at least 3 cm of ETT free length after strapping is applied
 - Recommended length of the ETT in ICU is at least 2 cm of ETT free length after ETT position is confirmed on CXR, and after strapping is applied.
3. Strapping of the ETT

Melbourne strapping is recommended for securing the ETT⁷. Follow the CoMET (East Midlands Children's Acute Transport) guideline for securing ETT.
4. Changing of the ETT strapping

A bolus of sedation together with a bolus of muscle relaxant is highly recommended. Cuff should always be deflated prior to extubation or re-positioning of ETT

Sedation

1. Sedation assessment is recommended at least 4th hourly utilizing COMFORT scale. (target COMFORT score between 17 and 26). Follow the sedation guidelines.

2. A bolus of sedation, with or without a bolus relaxant is suggested if required during tube suctioning, physiotherapy, patient hygiene, positioning, transport, and any other manipulation of the patient
3. All patients who require mechanical ventilation during intra-hospital transfer should be sedated and muscle relaxed. Follow CoMET guidelines for further details.

Outcome

1. Continuous UE event monitoring with regular at least quarterly review
 - Risk for UE in PICU/CICU is targeted to be lower than 1 per 100 ventilator days
2. Surrogate marker of monthly Airway Bundle Audit
 - Target 100% compliance with interim marker of 75%
 - A patient without all bundle elements would score as 'non-compliant' since you need all elements to achieve compliance
 - Aim to monitor at least 10 patients per unit per month

Review

This guideline should be reviewed after 3 years from date of approval and following results of clinical audit and future scientific evidence. The CPM lead retains responsibility for ensuring that review takes place in partnership with the PICU/CICU.

5. Education and Training

None

6. Monitoring Compliance

None currently identified

What will be measured to monitor compliance	How will compliance be monitored	Monitoring Lead	Frequency	Reporting arrangements

7. Supporting Reference

1. Sadowski R, Dechert RE, Bandy KP, Juno J, Bhatt-Mehta V, Custer JR, Moler FW, Bratton SL. Continuous quality improvement: reducing unplanned extubations in a pediatric intensive care unit. *Pediatrics* 2004 Sep;114(3):628-32
2. Lucas da Silva PS, de Carvalho WB. Unplanned extubation in pediatric critically ill patients: a systemic review and best practice recommendations. *Pediatr Crit Care Med*. 2010 Mar; 11(2):287-94
3. Rameshwar Mhamane, Nandini Dave and Madhu Garasia. Use of microcuff endotracheal tubes in paediatric laparoscopic surgeries. *Indian J Anaesth*, 2015 Feb; 59(2): 85-88
4. Dullenkopf A, Gerber AC, Weiss M. Fit and seal characteristics of a new paediatric tracheal tube with high volume-low pressure polyurethane cuff. *Acta Anaesthesiol Scand* 2005 Feb; 49(2):232-7
5. Schmolzer GM, Roehr CC. Techniques to ascertain correct endotracheal tube placement in neonates. *Cochrane Database Syst Rev*. 2014 Sep 13;9
6. Carvalho FL, Mezzacappa MA, Calil R, Machado Hda C. Incidence and risk factors of accidental extubation in a neonate intensive care unit. *J Pediatr (Rio J)*. 2010 May-Jun; 86(3):189-95
7. <https://resus.me/securing-infant-tracheal-tubes>
8. Capnography guidelines.pdf
9. Paulo Sergio Lucas da Silva; Maria Eunice Reis, Vânia Euzébio Aguiar and Marcelo Cunio, Machado Fonseca. Unplanned Extubation in Neonatal Intensive Care Unit: A Systematic Review, Critical Appraisal and Evidence-Based Recommendations. *RESPIRATORY CARE* December 27, 2012 as DOI: 10.4187/respcare.02164

10. Sandeep Tripathi, Denise J Nunez, Chaavi Katyal, and H Michael Ushay. Plan to Have No Unplanned: A Collaborative, Hospital-Based Quality-Improvement Project to Reduce the Rate of Unplanned Extubations in the Pediatric ICU. RESPIRATORY CARE Paper in Press. Published on May 19, 2015 as DOI: 10.4187/respcare.03984
11. D. Crankshaw, J. McViety, M. Entwistle. A review of cuffed vs uncuffed endotracheal tubes in children. *Pediatric Anesthesia and Critical Care Journal* 2014; 2(2):70-73
12. Weiss M, Balmer C, Dullenkopf A, et al. Intubation depth markings allow an improved positioning of endotracheal tubes in children. *Can J Anaesth* 2005; 52:721–726.
13. Dullenkopf A, Gerber Weiss M. Fit and seal characteristics of a new paediatric tracheal tube with high volume-low pressure polyurethane cuff. *Anaesthesiol Scand* 2005; 49:232–237.
14. De Orange FA, Andrade RGAC, Lemos A, Borges PSGN, Figueiroa JN, Kovatsis PG. Cuffed versus uncuffed endotracheal tubes for general anaesthesia in children aged eight years and under. *Cochrane Database of Systematic Reviews* 2017, Issue 11.
15. L Ford, J Ballard, M Davidson. Paediatric Intensive Care Unit (PICU) Guideline on the use of Cuffed Endotracheal Tubes Cuffed Endotracheal. *Glasgow Hospital Pediatric Intensive Care*
16. Josef Holzki, Michael Laschat, Christian Puder. Stridor is not a scientifically valid outcome measure for assessing airway injury. *Pediatric Anesthesia* 2009 19 (Suppl. 1): 180–197
17. Behzad Maghsoudi, Shojaolhagh Tariq, Seyed Basir Hashemi, Masoud Kaviani. The effect of cuffed endotracheal tube on respiratory complication following adenotonsillectomy in children. *Medical Journal of the Islamic Republic of Iran*. Vol. 24, No. 3, November, 2010. pp. 151-158
18. Timothy W. Deakers, George Reynolds, Mark Stretton, Christopher J.L. Newth. Cuffed endotracheal tubes in pediatric intensive care. *The Journal of Pediatrics* July 1994 Volume 125, Issue 1, Pages 57–62
19. C S Ashtekar, A Wardhaugh. Does cuffed endotracheal tubes increase the risk of airway mucosal injury and post-extubation stridor in children? *Best Evidence Topics*
20. Connie Taylor, Lekha Subaiya Daniel Corsino. Pediatric Cuffed Endotracheal Tubes: An Evolution of Care. *The Ochsner Journal* 11:52–56, 2011
21. Rennie C, Keightley J, Davidson M, *et al*
22. G368(P) An audit of unplanned extubations in a paediatric intensive care unit *Archives of Disease in Childhood* 2019;**104**:A150
23. PICANet annual report 2021
24. da Silva, P.S.L., et al., Care bundles to reduce unplanned extubation in critically ill children: a systematic review, critical appraisal and meta-analysis. *Archives of disease in childhood*, 2021. 12.
25. Kanthimathinathan, H.K., et al., Unplanned extubation in a paediatric intensive care unit: prospective cohort study. *Intensive Care Med*, 2015. 41(7): p. 1299-306.
26. The number of tracheal intubation attempts matters! A prospective multiinstitutional pediatric observational study: Lee et al. *BMC Pediatrics* (2016) 16:58 DOI 10.1186/s12887-016-0593-y
27. Razavi SS, Nejad RA, Mohajerani SA, Talebian M. Risk factors of unplanned extubation in pediatric intensive care unit. *Tanaffos*. 2013;12(3):11-6. PMID: 25191468; PMCID: PMC4153254.
28. Torres, Sherman J.; Palma, Carmela; and Mongiello, Michael, "Reducing Unplanned Extubations in the Pediatric Intensive Care Unit" (2020). Doctor of Nursing Practice (DNP) Projects. 26
29. da Silva, P. S.L., & Fonseca, M. C. M. (2017). Incidence and risk factors for cardiovascular collapse after unplanned extubations in the pediatric ICU. *Respiratory Care*, 62, 896-903. doi:10.4187/respcare.05346
30. Liang Chen, Jun Zhang, Guoshi Pan, Xia Li, Tianwu Shi, Wensheng He, Cuffed versus uncuffed endotracheal tubes in pediatrics: a meta-analysis. *Open Med*. 2018; 13: 366-373
31. Litman RS, Weissend EE, Shibata D, Westesson PL. Developmental changes of laryngeal dimensions in unparalyzed, sedated children. *Anesthesiology*. 2003 Jan;98(1):41-45.
32. Wani TM, Bissonnette B, Rafiq Malik M, Hayes D Jr, Ramesh AS, Al Sohaibani M, Tobias JD. Age-based analysis of pediatric upper airway dimensions using computed tomography imaging. *Pediatr Pulmonol*. 2016 Mar;51(3):267-71.
33. Jugpal TS, Garg A, Sethi GR, Daga MK, Kumar J. Multi-detector computed tomography imaging of large airway pathology: A pictorial review. *World J Radiol*. 2015 Dec 28;7(12):459-74.
34. Mizuguchi S, Motomura Y, Maki J, Baba R, Ichimiya Y, Tokuda K, Kaku N, Takada H, Maehara Y, Ohga S. Tracheal Size and Morphology on the Reconstructed CT Imaging. *Pediatr Crit Care Med*. 2019 Aug;20(8):e366-71.
35. Taylor C, Subaiya L, Corsino D. Pediatric cuffed endotracheal tubes: an evolution of care. *Ochsner J*. 2011 Spring;11(1):52-56.

36. Thomas R, Rao S, Minutillo C. Cuffed endotracheal tubes for neonates and young infants: a comprehensive review. *Arch Dis Child Fetal Neonatal Ed.* 2016 Mar;101(2):F168-74.
37. Chambers NA, Ramgolam A, Sommerfield D, Zhang G, Ledowski T, Thurm M, Lethbridge M, Hegarty M, von Ungern-Sternberg BS. Cuffed vs. uncuffed tracheal tubes in children: a randomised controlled trial comparing leak, tidal volume and complications. *Anaesthesia.* 2018 Feb;73(2):160-68.
38. de Wit M, Peelen LM, van Wolfswinkel L, de Graaff JC. The incidence of postoperative respiratory complications: A retrospective analysis of cuffed vs uncuffed tracheal tubes in children 0-7 years of age. *Paediatr Anaesth.* 2018 Mar;28(3):210-17
39. Mahmoud RA, Proquitté H, Fawzy N, Bühler C, Schmalisch G. Tracheal tube airleak in clinical practice and impact on tidal volume measurement in ventilated neonates. *Pediatr Crit Care Med.* 2011 Mar;12(2):197- 202
40. Salgo B, Schmitz A, Henze G, Stutz K, Dullenkopf A, Neff S, Gerber AC, Weiss M. Evaluation of a new recommendation for improved cuffed tracheal tube size selection in infants and small children. *Acta Anaesthesiol Scand.* 2006 May;50(5):557-61.
41. Effectiveness of Continuous Endotracheal Cuff Pressure Control for the Prevention of Ventilator-Associated Respiratory Infections: An Open-Label Randomized, Controlled Trial. Vu Quoc Dat,1 et al, *Clinical Infectious Diseases*, Volume 74, Issue 10, 15 May 2022, Pages 1795–1803

8. Key Words

Endotracheal tube, Melbourne strapping, Microcuff, Postextubation stridor, Subglottic stenosis

The Trust recognizes the diversity of the local community it serves. Our aim therefore is to provide a safe environment free from discrimination and treat all individuals fairly with dignity and appropriately according to their needs.

As part of its development, this policy and its impact on equality have been reviewed and no detriment was identified.

CONTACT AND REVIEW DETAILS

Guideline Lead (Name and Title)
Bedangshu Saikia - Consultant
Anand Patil – PICU Consultant

Executive Lead
Chief Nurse

Details of Changes made during review:

Pre-Intubation “Aide memoire”

Should be read aloud when team is ready

Prepare Equipment

Monitoring:

ECG (pulse tone on)

Spo2

BP (cycle every 2 min if no arterial line)

Capnography/Color capnometer

Airway items:

T-piece with mask

Suction on with Yankauer

Guedel/LMA

Laryngoscope

ET tubes (correct size & 1 size below and test the cuff)

Note ETT insertion length

ET Tapes ready

Stylet/Bougie

Magill forceps if nasal intubation

Drugs:

Induction agents & muscle relaxant dose confirmed

(decrease dose if hypotension likely)

Emergency drugs required? (Fluid bolus, Adrenaline)

Prepare Patient

*History reviewed? (Previous grade of laryngoscopy, past anaesthetic problems, previous ETT size/length)

*Airway assessment (any features of a predictably difficult airway)

*Adequately fasted? (Consider risks/benefits of delaying intubation/RSI if not fasted)

* Check access

*Pre-oxygenate for 2-3 mins with 100% O₂ (unless contraindicated)

* Optimise Position (Consider shoulder roll in <1 yr, pillow below head in >8 yr)

*Reliable access? (consider IO if difficult)

*Hemodynamics adequate (consider fluid bolus+/-peripheral vasoactive drugs pre-intubation)

*Aspirate NG tube

*Stethoscope

* Family aware of the plan?

Prepare Team

Assign roles

*Team leader

*1st Intubator

*2nd Intubator

*Intubator’s assistant

*Drug administrator

*NGT aspiration (continuously during facemask ventilation)

*Cricoid pressure

*Are we in best location? (move to PICU/theatre)

*Nurse in-charge aware?

*Consultant aware?

Prepare for failure

*Failure to intubate, CVS instability (who will deal and what should be done)

*Any specialist team required? (Anesthetist, ENT)

Confirm plan

Plan A: 2 attempts by 1st intubator

Plan B: Next attempt by senior

Plan C: Bag Mask Ventilation

Plan D: Call Anesthetist/ENT



Size of LMA		Size of ET suction catheter	
		2 times ETT size	
LMA size	weight (kg)	Intubation drugs (consider lower doses if risk of CVS instability)	
1	<5		
1.5	5-10		
2	10-20		
2.5	20-30		
3	30-50	Drugs	Dose
4	50-70	Ketamine	1-2mg/kg
		Fentanyl	1-2microgram/kg
		Rocuronium	1 mg/kg
		Atracurium	1 mg/kg

Sizes of ETT and insertion depth			
Age	Uncuffed ETT (mm ID)	Cuffed ETT (mm ID)	Length at lips (cm)
Neonate	3-3.5	3	9-10
>28 days 6 months	3.5	3-3.5	9-10
6 months-1year	3.5-4	3.5	10-12
2 years	4-4.5	4	12-13
5 years	5	4.5	14-15
8 years	5.5-6	5-5.5	16-18
10 years	6-6.5	6	17-18
12 years	6.5-7	6-6.5	18-20
15 years	7	6.5-7	20-21
18 years	7	7	20-21

Note: Rough guide

Size
 Uncuffed (> 1 year): Age /4 +4
 Cuffed (>3 kg): Age /4+ 3.5
 Neonate (under 3 kg): 2, 2.5 or 3.5 (Uncuffed)

Depth of ETT insertion
 Neonate: 3 times ETT size
 Oral: Age in yrs/2+ 12
 Nasal: Age in yrs/2 + 15 (older kids)
 Nasal: Age/2 + 14 (infancy and younger kids)