

## **1. Introduction**

- Hypocalcaemia is defined as plasma calcium < 2.1 mmol/L
- The commonest cause is post-surgical hypoparathyroidism after thyroidectomy
- Early symptoms include peri-oral and digital paraesthesia and carpopedal spasm
- Acute severe hypocalcaemia may lead to arrhythmias, seizures and acute laryngospasm
- ECG changes include prolonged QT interval and arrhythmias
- Hypocalcaemia associated with CKD should be discussed with the on-call renal physician
- The full list of causes should be considered:

### **Causes of hypocalcaemia**

- Post-thyroidectomy
- Post-parathyroidectomy
- Severe vitamin D deficiency
- Chronic Kidney Disease
- Magnesium deficiency
- Cytotoxic agents
- Rhabdomyolysis
- Pancreatitis
- Large volume transfusions

## **2. Scope**

This guideline applies to all staff when they are investigating and managing hypocalcaemia in an adult patient.

## **3. Recommendations, Standards and Procedural Statements**

- a) PTH, phosphate, ALP, U&E, Vitamin D and magnesium should be measured in all patients
- b) In acute severe hypocalcaemia, intravenous 10% calcium gluconate should be given (calcium chloride may also be used)
- c) Always consider and reverse the underlying cause of hypocalcaemia where possible
- d) See appropriate guidelines for severe Vitamin D deficiency and hypomagnesaemia where this is cause of hypocalcaemia

### **3.1 Management**

The management of hypocalcaemia should primarily be 'cause-specific'.

#### **Vitamin D deficiency**

Replace with loading or maintenance dose of colecalciferol as appropriate (see Leicestershire Medicines Strategy Group (LMSG) guidelines <http://www.lmsg.nhs.uk/?s=vitamin+d>)

#### **Post-thyroidectomy**

- a) Start oral calcium supplements (Calvive 1000 2 tablets bd or Adcal 3 tablets bd)
- b) In post-thyroidectomy patients repeat calcium in 24 hours
- c) Discharge to GP if calcium > 2.1 mmol/L
- d) If serum calcium remains low more than 72 hours post-operatively start 1-alfacalcidol 0.25mcg/day and refer for endocrinology or ENT clinic follow up as appropriate

## Hypomagnesaemia

- a) Stop precipitating drugs
- b) Consider other causes of hypomagnesaemia
- c) Give intravenous magnesium sulphate infusion (see UHL Hypomagnesaemia guidelines)  
[Hypomagnesaemia%20UHL%20Diabetes%20Guideline.pdf](#)

### **3.2 Severe hypocalcaemia (Serum calcium < 1.9 mmol/L or symptomatic below reference range)**

- a) This is a medical emergency – seek senior advice if patient unwell
- b) Ensure patient is on a cardiac monitor and in a clinically appropriate environment
- c) Administer 10mls 10% calcium gluconate in 50mls 5% dextrose over 10 minutes
- d) Repeat until patient asymptomatic to maximum of 40mls 10% calcium gluconate in 24 hours
- e) If no clinical response and airway is compromised then refer patient to ITU
- f) If patient stabilises and is asymptomatic give continuous calcium gluconate infusion (dilute 100ml 10% calcium gluconate in 5% dextrose or 0.9% sodium chloride and infuse at 50-100 ml / hour)
- g) Check calcium 4 hourly and titrate infusion until calcium normal and reversible cause treated
- h) In post-surgical hypoparathyroidism, start 1-alfacalcidol 0.25-0.5mcg/day and refer to endocrinology or ENT clinic as appropriate

### **Nursing interventions**

- a) Four hourly observations
- b) Temperature, pulse, BP, respirations, oxygen saturations
- c) More frequently if clinically indicated
- d) Patients with severe hypocalcaemia on IV infusion need continuous ECG monitoring
- e) Side effects of intravenous calcium include local thrombophlebitis, cardio-toxicity, hypotension, taste disturbance, nausea, flushing, vomiting and sweating

### **4. Education and Training**

None except dissemination of guideline

### **5. Monitoring and Audit Criteria**

Key performance indicators will be reduced morbidity and mortality caused by hypocalcaemia. Monitoring will be achieved by CMG mortality and morbidity reviews. Adverse incidents will be reported to CMGs.

### **6. Supporting Documents and Key References**

Society for Endocrinology Clinical Guidelines

### **7. Key Words**

Adult patient, management, hypocalcaemia

#### **CONTACT AND REVIEW DETAILS**

**Guideline Lead** Dr Miles Levy, Consultant Endocrinologist

**Trust Board Lead:** Medical Director

**Details of Changes made during review:** None